



Hygienic Considerations of Pathogenic *Escherichia Coli* Contamination on Cattle Carcass Surfaces in Egypt

Saad Mahmoud Saad¹, Amani Mohamed Salem¹, Shaimaa Nada², Mohamed – Mona²

¹ Food Control Department. Faculty of Veterinary Medicine, Benha University

² Animal Health Research Institute "Shebine EL-Koom Branch".

ABSTRACT

A total of 100 random samples were collected from surfaces of cattle carcasses in EL-Shouhada and Birket El –Sabaa Abattoirs in Menofya governorate (50 of each). The results showed that the mean *E. coli* counts were 5.44×10^2 and 7.99×10^2 cfu/cm² in EL- Shouhada and Birket EL-Sabaa abattoirs, respectively. Also, the incidence of *E. coli* isolated from cattle carcass surfaces in EL-Shouhada and Birket El-Sabaa abattoirs were 26% and 28%, respectively. The isolated serotypes of *E. coli* were EHEC as O₁₁₁ : H₂ (8%) and O₂₆ : H₁₁ (2%), EPEC as O₁₅ : H₄ (2%), O₅₅ : H₇ (2%), O₁₁₄ : H₄ (4%) and O₁₄₆ : H₂₁ (2%), ETEC as O₁₂₈ : H₂ (2%) and O₁₂₅ : H₂₁ (2%) and EIEC as O₁₂₄ (2%) in EL-Shouhada abattoirs, while in Birket EL-Sabaa abattoir were EHEC as O₁₁₁ : H₂ (6%) , O₂₆ : H₁₁ (8%) and O₁₀₃ : H₄ (2%), EPEC as O₅₅ : H₇ (2%) and O₉₁ : H₂₁ (2%) ETEC as O₁₂₈ : H₂ (6%) and EIEC as O₁₂₄ (2%). Findings of multiplex PCR showed that *eaeA* (intimin) gene was detected in (O₉₁:H₂₁ and O₁₁₄:H₄), *hlyA* (haemolysin) gene was detected in (O₁₂₅:H₂₁), *stx1* gene was not detected in the all isolated *E. coli* serogroups, but *stx2* gene was detected in (O₂₆:H₁₁ and O₁₁₄:H₄).

Keywords: *Escherichia Coli*, Cattle Carcass, Menofya governorate.

(<http://www.bvmj.bu.edu.eg>)

(BVMJ-34(1): 305-313, 2018)

1. INTRODUCTION

The sanitary conditions of abattoirs and their surrounding environment are major factors contributing in bacterial contamination of meat (Gill *et al.*, 2000).

Escherichia coli infection is highly prevalent in abattoir environment, and can pose a major threat to human health in underdeveloped communities (Abu El naga-*et al.*, 2014). Meanwhile, it is commonly non virulent but some strains have adapted pathogenic or toxigenic virulence factors that make them virulent for humans and animals (Malik and Memona, 2010). It is also

commonly used as surrogate indicator; its presence in food generally indicates direct and indirect fecal contamination (Clarence *et al.*, 2009).

Shiga-toxin producing *E. coli* (STEC) were found in a wide variety of animal species, including cattle, sheep, goats, pigs, water buffalos and wild ruminants (Caprioli *et al.*, 2005). Cattle form the main reservoir of (STEC) and fecal contamination of food represents the usual source of infection for humans but due to an apparently low infectious dose, human to human transmission

was also observed in outbreaks (Kuhnert *et al.*, 2000). Pathogenic strains of *E. coli* were divided into intestinal pathogenic *E. coli* (INPEC) causing diarrhea and extraintestinal pathogenic *E. coli* (EXPEC) including urinary tract infection (UTI), meningitis and septicemia depend on virulence factors and clinical symptoms (Kaper *et al.*, 2004 and Eid and Erfan, 2013).

Polymerase Chain Reaction (PCR) based methods were identified as a powerful diagnostic tool for the detection of pathogenic microorganisms (Malorny *et al.*, 2003). Compared to other methods of detection, these methods were rapid, highly specific and sensitive in the identification of target organisms (Wang *et al.*, 2007).

2. Materials and methods

2.1. Collection of samples:

A total of 100 random samples were collected from surfaces of cattle carcasses slaughtered in EL-Shouhada and Birket El – Sabaa abattoirs (50 of each) in Menofya governorate. The samples were taken using swab technique under aseptic conditions, just after washing and before stamping. The collected samples (Swabs) were preserved in an ice box then transferred to laboratory without undue delay and subjected to the microbiological examination.

2.2. Preparation of templates and Swabbing (FSIS, USDA., 1996)

A templates made of metal having an exposed inner area 10 cm^2 ($2 \times 5\text{cm}$) was used to delineate area of sampling.

Swabs from cattle carcasses surfaces were taken by using sterile cotton swabs and templates. The sterilized templates were placed firmly against the surface of examination to limit the examined area. The sterile cotton swab was rolled over the limited area inside the template, rolled in one direction and perpendicular to this direction to

represent all the examined area. Finally, the cotton swabs were aseptically retained into the rinsing fluid screw capped tubes containing 10ml buffered peptone water (0.1%).

2.3. Preparation of swabs (APHA, 2001):

The collected swabs were mixed in 225ml of sterile buffered peptone water (0.1%) to give 1/10 dilution. One ml from the original dilution was transferred with sterile pipette to another sterile test tube containing 9ml of buffered peptone water and mixed well to make the next dilution, from which further decimal serial dilutions were prepared.

2.4. Detection of *E. coli* count was done according to McFadden (2000).

2.5. Isolation and identification of *Escherichia coli* was done according to McFadden (2000).

2.6. Serological identification of *E. coli*

The isolates were serologically identified according to Kok *et al.* (1996) by using rapid diagnostic *E. coli* antisera sets (DENKA SEIKEN Co., Japan).

2.7. Polymerase Chain Reaction (PCR):

2.7.1. Extraction of DNA

2.7.2. Preparation of PCR Master Mix

2.7.3. Cycling conditions of the primers during cPCR

2.7.4. DNA Molecular weight marker

2.7.5 Agarose gel electrophoreses

3. RESULTS

The results in table (1) revealed that the mean of *E. coli* counts were $5.44 \times 10^2 \pm 1.13^a$ and $7.99 \times 10^2 \pm 1.13^A$ in EL-Shouhada and Birket El-Sabaa abattoir, respectively. On the other hand, the results in Table (3) showed that the incidence of *E. coli* in the examined swab samples was (26%) and (28%) in EL-Shouhada and Birket EL-Sabaa abattoirs, respectively. Also, it is obvious from the results achieved in Table (5) that the serological identified *E. coli* serotypes were EHEC as O₁₁₁ : H₂ (8%) and O₂₆ : H₁₁ (2%),

Hygienic Considerations of Pathogenic *Escherichia Coli* Contamination on Cattle Carcass Surfaces in Egypt

EPEC as O₁₅ : H₄ (2%), O₅₅ : H₇ (2%), O₁₁₄ : H₄ (4%) and O₁₄₆ : H₂₁(2%), ETEC as O₁₂₈ : H₂ (2%) and O₁₂₅: H₂₁ (2%) and EIEC as O₁₂₄ (2%) in EL-Shouhada abattoir. Meanwhile in Birket El-Sabaa abattoir, EHEC as O₁₁₁ : H₂ (6%) , O₂₆ : H₁₁ (8%) and O₁₀₃ : H₄ (2%), EPEC as O₅₅ : H₇ (2%) and O₉₁ : H₂₁(2%) ETEC as O₁₂₈: H₂ (6%) and EIEC as O₁₂₄ (2%). While, the results in Table (6) showed that the *eaeA* (intimin or *E. coli* attaching and effacing) gene was detected in (O₉₁:H₂₁ and O₁₁₄:H₄), *hlyA* (haemolysin) gene was detected in (O₁₂₅:H₂₁), *stx₁* gene was not detected in the all isolated *E. coli* serogroups, but *stx₂* gene was detected in (O₂₆:H₁₁ and O₁₁₄:H₄

Table (1): Mean values of *E. coli* count (cfu/cm²) in swab of cattle carcass surfaces from abattoirs in Menofyia governorate (n=50).

| Abattoirs | Min. | Max. | Mean± S.E* |
|-----------------|----------------------|----------------------|---|
| EL- Shouhada | 1.82×10 ² | 4.07×10 ³ | 5.44×10 ² ±1.13 ^a |
| Birket El-Sabaa | 2.29×10 ² | 6.31×10 ³ | 7.99×10 ² ±1.13 ^A |

S.E* = Standard error of mean

Table (2): Analysis of variance (ANOVA) of *E. coli* count in swab of cattle carcass surfaces from abattoirs in Menofyia governorate (n=50).

| Source of Variance | S.S. | D.F. | M.S. | F. value | Sig. |
|--------------------|--------|------|------|----------|------|
| Between Groups | .694 | 1 | .694 | 4.729 | .032 |
| Within Groups | 14.380 | 98 | .147 | | |
| Total | 15.074 | 99 | | | |

D.F = Degrees of freedom, S.S = Sum squares, M.S = Mean squares, = significant differences (P<0.05)

Table (3): Incidence of *E. coli* in the examined swab of cattle carcass surfaces from abattoirs in Menofyia governorate (n=50).

| Abattoir | Positive samples | |
|----------------|------------------|----|
| | NO. | % |
| EL- Shouhada | 13 | 26 |
| Birket El-Saba | 14 | 28 |
| Total | 27 | 54 |

Table (4): Serotypes of *E. coli* in the examined swab of cattle carcass surfaces from abattoirs in Menofya governorate (n=50).

| Swabs Strains | EL- Shouhada abattoir | | Birket-El-Sabaa abattoir | | Strain Characteristics |
|------------------|-----------------------|----|-----------------------------|----|------------------------|
| | No. | % | No. | % | |
| O111 : H2 | 4 | 8 | 3 | 6 | |
| O26 : H11 | 1 | 2 | 4 | 8 | EHEC |
| O103 : H4 | - | - | 1 | 2 | |
| O15 : H4 | 1 | 2 | - | - | |
| O55 : H7 | 1 | 2 | 1 | 2 | |
| O114 : H4 | 2 | 4 | - | - | |
| O146 : H21 | 1 | 2 | - | - | EPEC |
| O91 : H21 | - | - | 1 | 2 | |
| O128 : H2 | 1 | 2 | 3 | 6 | |
| O125 : H21 | 1 | 2 | - | - | ETEC |
| O124 | 1 | 2 | 1 | 2 | EIEC |
| Total | 13 | 26 | 14 | 28 | |

Table (5): The results of PCR amplifications of different used genes of *E. coli* serogroups:

| Sample | Results | | | |
|--|-------------|-------------|-------------|-------------|
| | <i>EaeA</i> | <i>HlyA</i> | <i>Stx1</i> | <i>Stx2</i> |
| 1(O ₁₅ :H ₄) | - | - | - | - |
| 2(O ₂₆ :H ₁₁) | - | - | - | + |
| 3(O ₅₅ :H ₇) | - | - | - | - |
| 4(O ₉₁ :H ₂₁) | + | - | - | - |
| 5(O ₁₀₃ :H ₄) | - | - | - | - |
| 6(O ₁₁₁ :H ₂) | - | - | - | - |
| 7(O ₁₁₄ :H ₄) | + | - | - | + |
| 8(O ₁₂₄) | - | - | - | - |
| 9(O ₁₂₅ :H ₂₁) | - | + | - | - |
| 10(O ₁₂₈ :H ₂) | - | - | - | - |
| 11(O ₁₄₆ :H ₂₁) | - | - | - | - |




Figure (1): Agarose gel electrophoresis of PCR amplification products using specific primers of (*eaeA*) gene of *E. coli*. Lane L: 100-600bp DNA Ladder. Pos.: control Positive at (248bp). Neg.: control Negative. Lane 1,2,3,5,6,8,9,10,11(Negative). Lane 4, 7 (Positive).




Figure (2): Agarose gel electrophoresis of PCR amplification products using specific primers of (*stx1*) gene, (*stx2*) gene and (*hlyA*) gene of *E.coli*.

**hlyA* : Haemolysin gene. Lane L: 100-1500bp DNA Ladder. Pos.: control Positive (at 1177bp). Neg.: control Negative. Lane 9 (Positive). Lane 1,2,3,4,5,6,7,8,10,11 (Negative).
 **Stx1*: Shiga toxin1 gene.Pos.: control Positive(at 614bp). Lane 1,2,3,4,5,6,7,8,9,10,11(Negative).

**stx2*: Shiga toxin 2 gene. Pos.: control Positive (at 779bp).

Lane 1,3,4,5,6,8,9,10,11(Negative). Lane 2, 7 (positive).

4. DISCUSSION

Foodborne illnesses caused by *E. coli* represent a major public health problem worldwide. Every treatment done to meat from the point of slaughtering until it is ready for consumption could add to the bacterial load of this meat. Thus, meat products are considered as a major vehicle of most reported foodborne outbreaks, and may be

contaminated with several types of organisms through long chain of preparation, handling of raw meat, processing, distribution, storage and retailing (Shawish, 2015).

The results recorded in Table (1) were similar with those reported by Bello *et al.*, (2011) (4.6 Log to 4.9 Log cfu/cm²), Ahmad (2013) (2.81 log cfu/cm²) and Eugène (2013) (0.8 to 3.0 log cfu/g). On the other hand, higher ones were recorded by EL-Morsi

(1994) (39×10^2 cfu/cm²) and Bogere, P. and Baluka, A. S. (2014) (8.4×10^4 cfu/g). While, lower ones obtained by Fliss *et al.* (1991) (0.15×10^2 organisms/cm²) and David *et al.* (2006) ($0.8 \log$ cfu/cm²). The results obtained in Table (3) nearly agreed with Moustafa (1993) (34%), Barkocy-Gallagher *et al.* (2003) (26.7%) and Abdelrahman- Alromisaa (2015) (25%). On the other hand, higher incidence were obtained by EL-Bassiouny and Samaha (1991) (80%), Nashid-Heba (1993) (48%), Enabulele and Uraih (2009) (100%) and Bogere and Baluka (2014) (83.3%). While, lower incidence was obtained by Hassouba (2000) (17.2%), Abdallah *et al.* (2009b) (8.86%). The variation in the results between counts and incidences of *E. coli* in the examined swab samples of cattle carcass surfaces between both of EL-Shouhada and Birket EL-Sabaa abattoirs was attributed to the differences in slaughtering, preparation, handling and the effectiveness of hygienic measures applied in these abattoirs. It is obvious from the results recorded in Table (5) that they were nearly similar to those obtained by Moustafa (1993) (O₂₆, O₅₅, O₁₁₁ and O₁₂₄), Abdelrahman- Alromisaa (2015) (O₁₂₈:H₂, O₁₂₄ and O₂₆:H₁₁) and Ismail-Eman (2015) (O₂₆:H₁₁ and O₁₂₈:H₂), while Barlow *et al.* (2006) could not isolated (O₂₆ and O₁₁₁), while Abdallah (2009) isolate (O₇₈ and O₈₆). The results in Table (7) and Fig (2) of PCR amplification of *Stx*₂ gene substantiated what was reported by Adrienne and James (1998), Ram *et al.* (2007) and Fernández *et al.* (2010). Meanwhile, the results in Table (7) and Fig (1) of PCR amplification of *intimin* gene were nearly agreed with those obtained by Adrienne and James (1998), Ram *et al.* (2007). Finally, PCR amplification of *HlyA* gene in examined isolates in Table (7) and Fig (2) substantiated what was reported by Adrienne and James (1998) and Wang *et al.* (2013).

As conclusion, the examined swab samples recovered from surfaces of carcasses slaughtered at both abattoirs were contaminated by pathogenic *E. coli*, but there was a highly incidence in samples which were collected from Birket EL-Sabaa abattoir than those from EL-Shouhada one. This conclusion was attributed to the differences in effective hygienic measures adopted in both abattoirs.

5. REFERENCES

- Abdallah, F. A. 2009. Prevalence of Enterobacteriaceae in cattle carcasses with special reference to pathogenic *Escherichia coli*. MVSc. thesis (Meat Hygiene), Fac. Vet. Med., Zagazig Univ., Egypt.
- Abdallah, M.A., Suliman, S.E., Ahmed, D.E., Bakhiet, A.O. 2009b. Estimation of bacterial contamination of indigenous bovine carcasses in Khartoum (Sudan). Afri. J. Microbiol. Res., 3(12): 882- 886.
- Abdelrahman-Alromisaa, A. M. E. 2015. Enteropathogenic *E. coli* in some slaughtered animals. M.V.Sc. Thesis (Meat Hygiene), Fac. Vet. Med., Zagazig, Univ., Egypt.
- Abu El Naga, Azza; S. M., Hedia, Riham H., Ata, Nagwa S., Zaki, Mona. S. 2014. Bacterial aspect of Food Poisoning. Life Sci. J., 11(3):290-298.
- Adrienne, W.P., James, C.P. 1998. Detection and characterization of shiga toxicogenic *Escherichia coli* by using multiplex PCR assay for *stx*₁, *stx*₂, *eae A*, *Enterohemorrhagic E. coli hlyA rfbO111* and *rfbO157*. J. of Clinical Microbiol., 36(2): 598-602.

- Ahmad, M. U. D., Safwat, A., Najeeb, M. I., Nawaz, M., Anjum, A. A., Ali, M. A., Mansur, N. 2013. Assessment of microbial load of raw meat at abattoirs and retail outlets. The J. of Animal and Plant Sciences, 23(3): 745-748 .
- American Public Health Association "APHA" 2001. Compendium of methods for microbiological examination of foods. 4th Edition 365-366-800. 1st, NW Washington DC 2000 1-3710.
- Banwart, G. J. (1981): Indicator organisms in: In: Basic food microbiology. 2nd ed. Avi. Publishing Co., Westport, Connecticut, USA.
- Barkocy-Gallagher, T. M., Arthur, M., Rivera-Betancourt, G. A., E. D. Berry, Nou, X., Shackelford, S. D., Wheeler, T. L. , Koochmaraie, M. 2003. Seasonal prevalence of Shiga toxin-producing *Escherichia coli* O₁₅₇:H₇ and non- 157 serotypes and Salmonella in commercial beef processing plants. J. Food Protect. 66(11): 1978.
- Barlow, R.S., Gpbious, K.S., Desmarchelier, P.M. 2006. Shiga toxin producing *Escherichia coli* in beef and lamb cuts. Inter. Food Microbiol. 11(1):105.
- Bello, M.; Lawan, K.; Kwaga, J. and Abiola, R. (2011): Assessment of carcass contamination with *E. coli* O157 before and after washing with water at abattoirs in Nigeria. International journal of Food Microbiology 150(2-3):184-186.
- Bogere, P. and Baluka, A. S. 2014. Microbiological Quality of Meat at the Abattoir and Butchery Levels in Kampala City, Uganda. Inter. J. Food Safety, (16): 29-35.
- Caprioli, A. S., Morabito, H., Bruge`re, E. Oswald. 2005. Enterohemorrhagic *Escherichia coli*: emerging issues on virulence and modes of transmission. Vet. Res. 36:289-311.
- Clarence, S.Y., Nwinyi, O., Chinedu, S.N. 2009. Assessment of bacteriological quality of ready to eat food (Meat pie) in Benin City metropolis, Nigeria. African J. of Microbiol. Research, 3 (6): 390-395.
- David, P., David, J., Stephan, M., Ian, J. , John, S. 2006. A National Survey of the Microbiological Quality of Beef Carcasses and Frozen Boneless Beef in Australia. J. Food Protect. 69(5): 1113.
- Egyptian Organization for Standardization and Specification "EOS". 2005. For Fresh meat.
- Eid, S.A.S., Erfan, A.M. 2013. Characterization of *E. coli* associated with high mortality of poultry flocks . Assiut Vet .Med. J. 59(139): 51-61, Egypt.
- El-Bassiouny, A., Samaha, H. 1991. Role of abattoir effluents in contaminating carcasses with some food poisoning bacteria. Assiut Vet. Med. J. 25 (49): 108, Egypt.
- El-Morsi, A. E. M. 1994. Sanitary status of buffalo-calf carcasses M.V.Sc. Thesis, Fac. Vet. Med., Zagazig Univ. Egypt.
- Enabulele, S.A., Uraih, N. 2009. Enterohemorrhagic *E. coli* O₁₅₇:H₇ Prevalence in meat and vegetables sold in Benin City, Nigeria. Afri. J. Microbiol. Res., 3(5): 276-279.
- Ercolini, D., Russo, F., Torrieri, E., Masi, P., Villani, F. 2006. Changes in the spoilage-related microbiota of beef

- during refrigerated storage under different packaging conditions. *Appl. Environ. Microbiol.* 72 (7): 4663-4671.
- Eugène, N.; Divine, B. and Martin, P. O. (2013): Assessment of beef meat microbial contamination during skinning, dressing, transportation and marketing at a commercial abattoir in Kigali city, Rwanda. *PAK. J. Food Sci.*, 23(3): 133-138.
- Fernández, D., Irino, K, Sanz, M. E., Padola, N. L., Parma, A. E. 2010. Characterization of Shiga toxin-producing *Escherichia coli* isolated from dairy cows in Argentina. *Lett. Appl. Microbiol.* 51(4):377-82.
- Fliss, I., Simard, R.E. , Etriki, A. 1991b. Microbiological quality of different fresh meat species in Tunisian slaughter houses and markets. *J. Food Protect.* 54:773-777.
- Food Safety and Inspection service “FSIS” United States Department of Agriculture .1996. Pathogen Reduction; Hazard Analysis and Critical Control Point (HACCP) Systems; Final Rule. Federal Register/ Vol. 61, No. 144/ Thursdat, July 25.
- Gill C. O., Bryant, J., Brereton, D. A. 2000. Microbiological conditions of sheep carcasses from conventional or inverted dressing processes. *J Food Protect.* 63(9): 1291-1294.
- Hassouba, M. M. 2000. Prevalence of enteric pathogens in meat. Ph. D. Thesis, Fac. Vet. Med., Cairo Univ., Egypt.
- Ismail- Eman, M. H. 2015. Improvement of Microbiological State of Menofia Abattoirs. M.V.Sc. Thesis (Meat Hygiene) Fac. Vet. Med. Moshtohor, Benha Univ.
- Kaper, J. B., Nataro, J. P., Harry, L.T. 2004. Pathogenic *Escherichia coli*. *Nat. Rev. Microbiol.* 2:123-140.
- Komba, E.V.G., Mkupasi, E. M., Mbyuzi, A. O., Mshamu, S., Luwumbra, D., Busagwe, Z. , Mzula, A. 2012. Sanitary practices and occurrence of zoonotic conditions in cattle at slaughter in Morogoro Municipality, Tanzania: implications for public health. *Tanzania J Health Res.* 14 (2): DOI: <http://dx.doi.org/10.4314/thrb.v14i2.6>.
- Kuhnert, P., Boerlin, P., Frey, J. 2000. Target genes for virulence assessment of *Escherichia coli* isolates from water, food and the environment. *FEMS Microbiol. Reviews* (24): 107-117.
- McFadden, J. F. 2000. Biochemical tests for identification medical bacteria. Waryery Press Inc., Baltimore, Md. 21202 USA.
- Malik, K., Memona, H. 2010. Molecular and immunological studies of pathogenic *Escherichia coli* in meat samples collected from different localities of Lahore. *International J. Cell and Molecular Biol. (IJCMB)*, 1 (3) 218-224.
- Malorny, B., Tassios, P.T., Radstrom, P., Cook, N., Wagner, M., Hoorfar, J. 2003. Standardization of diagnostic PCR for the detection of food borne pathogens. *Inter. J. Food Microbiol.* 1 (25): 39-48.
- Moustafa, M.M.M. 1993. Bacteriological studies on certain Gram negative food borne pathogens. Ph.D., Thesis, Military Institute of Health and Epidemiology, Military Medical Academy.
- Nashid- Heba, F. 1993. Salmonella and Enteropathogenic *E. coli* serotype in

meat and meat products. M.V.Sc. Thesis (Meat Hygiene) Fac. Vet. Med. Moshtohor, Zagazig Univ., Benha Branch, Egypt.

Ram, S., Vajpayee, P., Hanker, R. 2007. Prevalence of multi antimicrobial-agent resistant, shiga toxin and enterotoxin producing *Escherichia coli* in surface waters of River Ganga. Environmental Science and Technology 41: 7383-7388.

Shawish, R. R. M. 2015. "Prevalence of shiga toxin-producing *Escherichia coli* in some beef products". Ph.D. Thesis (Meat Hygiene), Fac. Vet. Med., Menoufia Univ. (Sadat Branch), Egypt.

Sousa, C. P. 2008. "The Impact of Food Manufacturing Practices on Food borne Diseases". Brazilian Archives of Biology and Technology 51(4), 815-823.

Tebbut, G. M. 1999: Microbiological contamination of cooked meats and environmental site in premises selling

both raw and cooked meat products. Inter. J. Environ. Health Res. 3(4): 209-216.

Wang, L., Li, Y., Mustapha, A. 2007. Rapid and simultaneous quantification of *Escherichia coli* O₁₅₇:H₇, salmonella and shigella in ground beef by multiplex real-time PCR and immunomagnetic separation. J. Food Protect. 70(6):1366-1372.

Wang, X. G., Zhang, Y. H., Chen, X. H., Luo, L. F., Liu, Y., Liu, J. Q., Song, C. P., Chen G. Q. 2013. Establishment and application of multiplex PCR for non-0157:H7 STEC virulence genes detection. Zhonghua Shi Yan He Lin Chuang Bing Du Xue Za Zhi. (Article in Chinese), 27(5):388-91.